

Solving Problems Using Problem Solving

Dr. David C. Stone
STAO Annual Conference, Toronto
November 11th 2011

Contact information:



Tel.: 416-946-0293

Email: dstone@chem.utoronto.ca

Web: <http://www.chem.utoronto.ca/~dstone/teachers/>

A big problem:

- There is **very little correlation** between high school grades and subsequent performance in university, even when the course content appears to be simple 'review' of the high school curriculum...
- 25% of students see their grade **drop by 30-60 percentage points** in first year undergraduate chemistry compared to high school, irrespective of the high school grade earned...
- The best studies to date can only explain <40% of the variation in student achievement in first year chemistry (or other science subjects)...
- Students who struggle lack confidence, are often weak in algebra, and tend to approach learning as an exercise in memorisation without understanding...



So what do the ones who struggle get wrong, why, and what can we do about it?

Think in terms of: Piaget (concrete versus formal development – most high school chemistry requires formal thought but most students have not developed that ability yet); Perry (intellectual development – students tend to be dualist black/white, right/wrong thinkers and often have difficulty correctly understanding models or reconciling multiple models); and Bloom's taxonomy (simplified – there's much that needs to be remembered at the basic level, but we need to push towards understanding concepts and applying them to new situations).

Self-Test/Assessment Questions

The following are all taken from 1st and 2nd-year university multiple-choice student self-assessment tests. The distractors in some of the questions were deliberately chosen based on known errors that students commit; this enables highly specific feedback to be provided to the students based on their choices for each question. Such an approach is not recommended for high-stakes tests or exams!

- Given the expression $A/B = X/Y$, then B is equal to:
 - XY/A
 - AY/X**
 - X/AY
 - Y/AX
- When correctly expressed in SI units, a density of 1.23 g/cm³ is:
 - 1.23 x 10⁻³ g/m³
 - 1.23 x 10⁻³ kg/m³
 - 1.23 g/m³
 - 1.23 x 10³ kg/m³**
- Consider the following balanced chemical reaction:
$$2 \text{MnO}_4^- + 16 \text{H}^+ + 15 \text{I}^- \rightarrow 2 \text{Mn}^{2+} + 5 \text{I}_3^- + 8 \text{H}_2\text{O}$$
What volume of 0.0525 M iodide would be required to exactly react with 20.0 ml of 0.0125 M permanganate?
 - 0.63 ml
 - 4.76 ml
 - 35.7 ml**
 - 84.0 ml
- A solution of known iodine concentration may be prepared by mixing solutions of iodate and iodide under acidic conditions:
$$a \text{IO}_3^- + b \text{I}^- + c \text{H}^+ \rightarrow p \text{I}_2 + q \text{H}_2\text{O}$$
When correctly balanced, the stoichiometric coefficients in this reaction equation are:
 - $a = 1, b = 1, c = 6, p = 1, q = 3$
 - $a = 1, b = 5, c = 6, p = 3, q = 3$**
 - $a = 3, b = 3, c = 6, p = 3, q = 3$
 - $a = 5, b = 1, c = 6, p = 1, q = 5$
- A 500 mL sample of a solution contains 0.375 moles of HNO₃ in pure water. What is its pH?
 - pH 0.125**
 - pH 0.426
 - pH 0.727
 - The pH cannot be less than 1

The following two questions appear back-to-back on the University of Guelph 1st-year chemistry student evaluation test. The results obtained are consistent year-over-year...

6. The geometry of a water molecule is:
 - a. angular or bent
 - b. linear
 - c. pyramidal
 - d. tetrahedral

7. The geometry of a molecule in which the central atom is bonded to 2 H atoms and has 2 lone pairs is:
 - a. angular or bent
 - b. linear
 - c. pyramidal
 - d. tetrahedral

Notes: question 1 is typical of every chemistry diagnostic exam or test of formal (logical) thinking development; the ability to do basic algebraic manipulations of equations – and especially those involving ratios – is essential to doing well in physical sciences. The following tables summarises student response rates for each option in the preceding multiple-choice questions. Correct answers are in bold font.

Question	a	b	c	d	Comment
1	7.9%	68.4%	18.4%	5.3%	
2	4.8%	28.6%	4.8%	61.9%	(b) inverted conversion factors
3	7.0%	18.6%	72.1%	2.3%	(a) inverted coefficients (b) omitted coefficients (d) inverted concentrations
4	51.2%	41.9%	7.0%	0%	(a) balanced mass only (c) balanced charges only
5	67.9%	10.7%	7.1%	11.9%	(b)(c) incorrect conc'n
6	88.8%	4.5%	4.6%	2.1%	
7	15.9%	53.0%	13.9%	17.2%	

Note: The terminology applied to the examples throughout this handout follows a simplified version of Bloom's taxonomy, consisting of **Recall**, **Algorithmic** (or **Algebraic**), and **Conceptual** questions. This provides a much easier means of checking whether or not a test or exam meets the instructor's intended goals as far as the types of evaluation. For a typical 1st- or 2nd-year undergraduate course, for example, the marks might be distributed amongst the three categories of question as 33% R, 33% A, and 33% C.

Troublesome Knowledge & Alternate Conceptions (Naive conceptions, misconceptions)

Troublesome Knowledge (Perkins):

“Some students will resort to rote memory and routine procedures ... They will try to learn enough ... to pass the test without developing any real insider feel. And pass they may, ending up with *knowledge troubled by partial and brittle understandings...*”

Alternate conceptions arise from (Kind, 2004; Talanquer, 2006; Taber & Garcia Franco, 2009):


- “Common sense” reasoning
- Intuitive thinking
- Personal experience
- Mental shortcuts (“cognitive misers” – Keith Stanovich, 2009)
- Misuse of analogies (inappropriate or limits not stated)
- Misunderstanding of the nature of scientific models
- The way we teach and grade assessments

“Many of students’ alternative conceptions in chemistry seem to result from the confident and impulsive application of a crude, incomplete, limited, and superficial explanatory framework about chemical substances and phenomena. This knowledge system ... creates the illusion of explanatory depth: **students believe that they understand more than they actually do.**”
(Talanquer, 2006, emphasis added)

Example 1:

- Balance the following equation for the production of ammonia: $N_2 + H_2 \rightarrow NH_3$
- Represent the balanced reaction using circles with letters in the centre to depict the atoms:

Here is what one student drew:



*A common misconception, possibly stemming from the way we describe splitting a substance into smaller and smaller units, is that individual atoms have the **same** physical properties as their macroscopic substance, e.g. Cu atoms have the same properties as bulk metallic copper; this is plainly false! The following questions are adapted from Ref. 1:*

Example 2:

A metallic wire has the following properties: a) conducts electricity; b) brown colour; c) density of 8.93 g/cm^3 ; d) malleable and ductile; e) expands on heating. Suppose you could isolate one single atom from the metallic wire: which of the above properties would it have?

None of the above! For density, remember that there are voids between the atoms even with the closest packing arrangement.

The wire is heated in an evacuated vessel until it completely evaporates. The resulting gas has the following properties: a) compressible; b) expands on heating; c) pungent odour; d) yellow colour; e) attacks plastics. Suppose you could isolate one single atom from this gas: which of the above properties would it have?

*You could debate about c and e, until you remember that we are talking about **single** atom here...*

The following are all examples taken from concept tests. Note particularly the gas law example, which shows how you can use the same question to develop and subsequently test for specific concepts. In addition, there are many conceptual problems and a "Chemistry Concepts Inventory" (similar to the physics Force Concepts Inventory) available from the J. Chem. Ed. question bank site: <http://jchemed.chem.wisc.edu/JCEDLib/QBank/collection/index.html>

Mole Concept Test (From Duncan & Johnstone, *Education in Chemistry*, 1973, 213-214)

- Given that $\text{Mg} + \text{S} \rightarrow \text{MgS}$, what mass of Mg would react completely with 32 g of S?
 - 12 g
 - 24 g
 - 32 g
 - 56 g
- Given that $2 \text{NaOH} + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{SO}_4 + 2 \text{H}_2\text{O}$, how many moles of H_2SO_4 are required to react with 1 mole of NaOH?
 - $\frac{1}{2}$ mole
 - 1 mole
 - 2 moles
 - 4 moles
- Given that $2 \text{SO}_2 + \text{O}_2 \rightarrow 2 \text{SO}_3$, what mass of SO_2 would react with 32 g of O_2 ?
 - 32 g
 - 64 g
 - 96 g
 - 128 g
- Given that $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$, what mass of O_2 is required to react with 3 g of carbon?
 - 8 g
 - 16 g
 - 64 g
 - 128 g
- What is the correctly balanced form of the equation $\text{Al} + \text{O}_2 \rightarrow \text{Al}_2\text{O}_3$?
 - $2 \text{Al} + \text{O}_3 \rightarrow \text{Al}_2\text{O}_3$
 - $\text{Al}_2 + 3 \text{O} \rightarrow \text{Al}_2\text{O}_3$
 - $4 \text{Al} + 3 \text{O}_2 \rightarrow 2 \text{Al}_2\text{O}_3$
 - $\text{Al}_4 + 6 \text{O} \rightarrow 2 \text{Al}_2\text{O}_3$
- Given that $\text{N}_2 + \text{H}_2 \rightarrow \text{NH}_3$, how many moles of H_2 would react completely with 1 mole of N_2 ?
 - 1 mole
 - 2 moles
 - 3 moles
 - 4 moles
- A molar solution of HCl contains
 - 1 mole of HCl dissolved in 1 mole of water
 - 1 mole of HCl dissolved in 1 litre of water
 - 1 mole of HCl dissolved in 1 litre of solution
 - 1 mole of water dissolved in 1 litre of HCl

8. Which solution of NaCl is most concentrated?
- 200 mL of a solution containing 2 moles of dissolved NaCl
 - 500 mL of a solution containing 4 moles of dissolved NaCl
 - 750 mL of a solution containing 8 moles of dissolved NaCl
 - 1000 mL of a solution containing 6 moles of dissolved NaCl
9. Which of the following solutions of HCl is most concentrated?
- 500 mL of 2 M HCl
 - 1000 mL of 3 M HCl
 - 300 mL of 4 M HCl
 - 800 mL of 5 M HCl
10. Which of the following solutions contains the most NaCl?
- 500 mL of 2 M NaCl
 - 1000 mL of 3 M NaCl
 - 250 mL of 4 M NaCl
 - 200 mL of 5 M NaCl

Notes and comments on questions 1–10 from the original study of Scottish high school students: Facility value (FV) is the percentage answering the question correctly; discrimination power (DP) is the difference in FV values between the top and bottom third of the students when ranked by overall score (for large classes, upper and lower quartiles can be used instead).

Num.	Answer	FV(%)	DP(%)	Comment
1	b	80	42	
2	a	51	55	Deviation from 1:1 stoichiometry causes problems
3	d	37	57	Similar comment
4	a	72	41	
5	c	55	39	Balancing equations problematic
6	c	37	29	Almost half chose answer (a)
7	c	55	22	Students unaware of finite volume of <i>solute</i> in solvent!
8	c	>66%	n/a	
9	d	42	15	Low DP implies even best students making mistakes
10	b	39	10	Most ignored the volume in the problem

Additional Note:

Half of 2nd-year undergraduates in an analytical chemistry course incorrectly chose b for question 7 on a self-assessment test

Additional conceptual questions can be found on the J. Chem. Ed. Question Bank web site at:
<http://jchemed.chem.wisc.edu/JCEDLib/QBank/collection/CCInventory/index.html>

Structured, Diagnostic, and Conceptual questions

This question is an example of conceptual vs. algorithmic (“algebraic”) problem solving, and can be asked in one of three ways. Source: Lillian Bird, *J. Chem. Ed.*, **2010**, 87(5), 541-546.

Structured example:

A flask is first evacuated, then filled with 0.200 g of methane gas (CH₄) and stoppered.

- The molecular mass of methane is 16.0 g/mol. How many moles of methane are contained in the flask?
- The flask containing the methane is held at a constant temperature of 298 K. What is the pressure if the flask has a volume of 1.00 L? The value of the gas constant $R = 0.0821 \text{ atm L}/(\text{mol K})$.
- A second flask is filled with 0.200 g of carbon dioxide gas (CO₂, molecular mass of 44.0 g/mol). How many moles of carbon dioxide are contained in the second flask?
- The flask containing the carbon dioxide is also held at a constant temperature of 298 K. What is the pressure in this flask if it also has a volume of 1.00 L? The value of the gas constant $R = 0.0821 \text{ K}/(\text{mol L})$.
- Compare your answers to b) and d). How does the pressure in each flask relate to the molecular weight of the gas within it, given that both have the same volume and are at the same temperature?
- Use this information to predict whether a 1.00 L flask containing 0.200 g of nitrogen gas (N₂, molecular weight 28.0 g/mol) at 298 K would have a higher or lower pressure than the flask containing the carbon dioxide. Explain your reasoning.

Diagnostic example:

Individual 0.200 g samples of each of the following gases were placed in four separate 1.00 L stoppered flasks at 298 K. In which flask do you expect the gas to exert more pressure? Explain your answer.

Flask:	1	2	3	4
Gas in flask:	CH ₄	Ne	N ₂	CO ₂
M_m (g/mol)	16.0	20.2	28.0	44.0

Lillian Bird notes that of 106 students: 36% calculated the pressure for each gas and justified their answer solely on these values; 42% similarly did all the calculations but then derived the principle and used that as justification; 26% gave the correct answer solely in terms of principles.

Concept text example:

Four identical sealed containers are each filled with a different gas (as indicated below) until each contains exactly the same mass. If all four are held at the same temperature, which flask contains gas at the greatest pressure?


This can be provided with or without the same table as before; students can only provide a solution if they can manipulate the concepts and equations *without* the benefit of a concrete calculation, since the mass, volume, and temperature are never stated!

Conceptual *versus* Algorithmic (Algebraic) Thinking

Mary Nakhleh and others have adopted a diagnostic approach using *paired* questions: the first usually requires a fairly straight-forward calculation that can be done simply by following a memorized procedure (algorithmic or algebraic thinking); the second explores a student's *conceptual* understanding of the same phenomenon. A number of the alternate conceptions test questions use the same structure.


Gas laws example:

- 0.100 mole of hydrogen gas occupies 600 mL at 25 °C and 4.08 atm. If the volume is held constant, what will be the pressure of the sample of gas at -5 °C?
 - 2.98 atm
 - 3.67 atm
 - 4.08 atm
 - 4.54 atm
 - 6.00 atm
- The diagram on the left represents a cross-sectional view of a rigid sealed tank filled with hydrogen gas at 20 °C and 3 atm pressure. The dots represent the distribution of hydrogen molecules within the tank. Which of the diagrams (a) through (e) illustrates one possible distribution of the hydrogen gas molecules in the tank if its temperature is lowered to -5 °C? The normal boiling point of hydrogen is -252.8 °C.



Limiting reagent example (actually three related questions):

- Which is the limiting reagent when 2.0 mol of CO₂ reacts with 2.0 mol of S₂ to form COS and O₂?
 - CO₂
 - S₂
 - COS
 - O₂
- Atoms of three different elements are represented by ⊗, ⊕, and ∅. Which is the limiting reagent when two ⊗⊗ molecules and two ∅∅⊕ molecules react to form ⊗∅⊕ and ∅∅?
 - ⊗⊗
 - ∅∅⊕
 - ⊗∅⊕
 - ∅∅
- The diagram on the left represents a mixture of S atoms and O₂ molecules in a closed container. Which diagram on the right shows the results after the mixture reacts as completely as possible according to the equation 2S + 3O₂ → 2SO₃?



Other paired examples:

4. When a match burns, some matter is destroyed – True or False?
5. What is the reason for your answer to question 4?
 - a. This chemical reaction destroys matter.
 - b. Matter is consumed by the flame.
 - c. The mass of ash is less than the match it came from.
 - d. The atoms are not destroyed, they are only rearranged.
 - e. The match weighs less after burning.


6. A 1.0-gram sample of solid iodine is placed in a tube and the tube is sealed after all of the air is removed (see diagram). The tube and the solid iodine together weigh 27.0 grams. The tube is then heated until all of the iodine evaporates and the tube is filled with iodine gas. Will the weight after heating be:
 - a. Less than 26.0 grams.
 - b. 26.0 grams.
 - c. 27.0 grams.
 - d. 28.0 grams.
 - e. More than 28.0 grams.
7. What is the reason for your answer to question 6?
 - a. A gas weighs less than a solid.
 - b. Mass is conserved.
 - c. Iodine gas is less dense than solid iodine.
 - d. Gasses rise.
 - e. Iodine gas is lighter than air.

8. 100 mL of water at 25°C and 100 mL of alcohol at 25°C are both heated at the same rate under identical conditions. After 3 minutes the temperature of the alcohol is 50°C. Two minutes later the temperature of the water is 50°C. Which liquid received more heat as it warmed to 50°C?
 - a. The water.
 - b. The alcohol.
 - c. Both received the same amount of heat.
 - d. It is impossible to tell from the information given.
9. What is the reason for your answer to question 8?
 - a. Water has a higher boiling point than the alcohol.
 - b. Water takes longer to change its temperature than the alcohol.
 - c. Both increased their temperatures 25°C.
 - d. Alcohol has a lower density and vapor pressure.
 - e. Alcohol has a higher specific heat so it heats faster

Contributed Problems from Participants

(Partially re-written!)

1. Consider the graph below, which shows the relationship between temperature (T) and volume (V) for a gas. This relationship holds for all gases under all conditions – True or False?



What is the reason for your answer?

- (a) At very low temperatures, atoms have finite volume
- (b) Because it is always T/V
- (c) If the mass changes, V changes
- (d) Because it follows Charles' Law

Comments: This is a good start for a two-part question, especially as part of group discussions where students have to justify their answers to one another. I'm not sure if the choice of axes was intentional – T is the independent variable, so should be on the x-axis – but it does make for another interesting discussion point!

2. A flask contains 500 mL of a 6.0 mol/L solution of HCl. If 250 mL is used for a titration, the concentration of the solution remaining in the flask is
- (a) 3.0 mol/L
 - (b) 6.0 mol/L
 - (c) 9.0 mol/L
 - (d) 12.0 mol/L

What is the reason for your answer?

- (a) Concentration does not change
- (b) Half the liquid was removed so the concentration increases
- (c) Half the volume means half the concentration

Comments: This is similar to one of the chemistry concept inventory questions, in which a mass of NaCl is added to a saturated solution of NaCl.

3. Consider the molecules methane (CH_4) and dichloromethane (CH_2Cl_2). In comparing these two molecules:
- (a) Both have the same polarity
 - (b) Methane is more polar than dichloromethane
 - (c) Dichloromethane is more polar than methane

What is the reason for your answer?

- (a) Both molecules are symmetrical
- (b) Neither molecule contains a highly electronegative atom
- (c) C–H bonds are more polar than C–Cl bonds
- (d) Tetrahedral molecules are always non-polar
- (e) Dichloromethane contains a mix of bond polarities

There was also an interesting mass conservation/buoyancy question, but I wasn't clear on the reaction involved and, given the rate at which students avoid physics, I felt the buoyancy component might actually cause more confusion.

It does, however, give me the opportunity to remind you that ALL balances – mechanical or electronic – actually determine WEIGHT and not mass. The only reason we can treat the weight as the mass is because they are calibrated so as to give the reading in mass units. This does, however, neglect the issue of buoyancy, so you need to make corrections if measuring the mass of gas in a balloon, for example.

Good Problems, Bad Problems

Sometimes, you can create problems where none existed. At other times, you can create opportunities for thought, discussion, and reflection. As an example, consider how *your* students would likely respond to the following two questions, posed in this exact sequence, on a test or exam:

1. What is the pH of an acid?
2. What is the pH of 1.0×10^{-8} mol/L of HCl?

Points raised during discussion:

- Most students would answer “pH 8” and not realise the mistake
- This pair of questions is good for small group discussion, but not so good for a high-stakes test or exam
- A good way to introduce the autoionization of water
- Supplementary questions such as “does your answer make sense?” and “what is the primary source of protons in the solution?” can be used to promote thought/discussion
- The calculation of actual pH for acids when the formal concentration of the acid is 10^{-6} M or less is possible but actually quite complicated – both mass (atom) and charge balances must be set up to solve the equilibrium expressions for both water and the acid since the normal simplifying assumptions in solving the ICE table equation cannot be made

Problem Solving

John Hayes: “Whenever there is a gap between where you are now and where you want to be, and *you don’t know how* to find a way to cross that gap, you have a problem” (emphasis added; as cited by Bodner, 2003).

Polya’s model of problem solving (ok for routine exercises but...) *C.f.* “GRASS”

- Understand the problem
- Devise a plan
- Carry out the plan
- Look back

The Anarchistic Route, or “How an *expert* solves problems” (Wheatley, as cited by Bodner):

- **Read the problem**
- Now read the problem again
- Write down what you hope is the relevant information
- **Draw a picture**, make a list, or write an equation or formula to help you begin to understand the problem
- Try something
- Try something else
- See where this gets you
- Read the problem again
- Try something else
- See where this gets you
- Test intermediate results to **see whether you are making any progress** toward an answer
- Read the problem again
- When appropriate, strike your forehead and say, “son of a...”
- Write down ‘an’ answer (not necessarily ‘the’ answer)
- **Test the answer to see if it makes sense**
- Start over if you have to, celebrate if you don’t

The Waterfall Problem

The Horseshoe Falls are 49 m high. Assuming that all the potential energy of the water is converted into heat, how much warmer is the water at the bottom of the falls than at the top? Comment on the magnitude of your answer. The flow of water over the falls is reduced at night as more is diverted through the hydroelectric generating station. What affect will this have? Give reasons for your answers.

Relevant Data:

Potential energy $E_p = mgh$ where m is mass (kg), $g = 9.81 \text{ m s}^{-2}$, and h is height (m); the specific heat capacity of water $s = 4.179 \text{ J/(g } ^\circ\text{C)}$.

Solution:

First, we need to know how much energy is available to be converted into heat. This is the potential energy associated with the water being at the top of falls. Since we don't know how much water is going over the falls, the simplest approach is to calculate for unit mass:

$$E_p = mgh = 1 \times 9.81 \times 49 \text{ kg} \cdot \text{m s}^{-2} \cdot \text{m} = 481 \text{ kg m}^2 \text{ s}^{-2} \text{ or } 481 \text{ J}$$

Next, we need to convert this to a change in temperature. Remember that *changes in state* are always (final – initial), so $\Delta T = (T_{\text{final}} - T_{\text{initial}})$. We assume that all the potential energy is converted into heat, so that:

$$E_p = q = sm\Delta T \therefore \Delta T = \frac{E_p}{sm} = \frac{481}{4.179 \times 1 \times 10^3} \frac{\text{J}}{\text{J g}^{-1} \text{ } ^\circ\text{C}^{-1} \cdot \text{kg} \cdot \text{g kg}^{-1}} = \mathbf{0.12 \text{ } ^\circ\text{C}}$$

This is a small but reasonable value, given the assumptions made. When the flow of water is reduced overnight, the potential energy of each kilogram of water at the top of the falls is still the same, so the warming will be the same. Heat transfer to the surroundings will obviously be different at night, but we did not consider that in our initial calculation either.

Comments:

Some students don't even know where to start this problem, as they are unused to connecting concepts from different parts of the course (let alone different courses!) Many make a common mistake with units and predict an extremely large temperature difference that is in excess of water's normal boiling point. Of these, some state that it "looks wrong" but many don't comment at all.

In discussing this question, point out that students need to identify the core concepts first, rather than simply 'plugging and playing' with the first equation that seems relevant. It is also very helpful to emphasize unit analysis for physical chemistry problems such as this, as errors can often be caught here. When presenting the solution, be sure to state the *thought process* at each step before the actual calculation! Finally, since the height is only given to 2 s.f., the final answer can only be stated to 2 significant figures.

The Pizza Problem

We all know that if you try and eat pizza too soon after it comes out of the oven you can burn your mouth. Is this because of the crust, the cheese, or the tomato sauce? Use your knowledge of the different phases of matter, kinetic molecular theory, and thermochemistry to justify your answer.

Some relevant data:

- Assume that a 50 g slice of pizza consists of 25 g crust, 20 g cheese, and 5 g of sauce. Approximate values of the specific heat capacity are: Cheese = 3.0 J/(g °C); Crust = 2.0 J/(g °C); Sauce = 4.0 J/(g °C). The heat capacity of water = 4.2 J/(g °C).
- Pizza is cooked at ~ 450 °F (230 °C). Assume that, by the time it reaches your table, the pizza has reached a uniform temperature of about 150 °C. The soft tissue on the inside of your mouth (which has a very high water content) is 37 °C. Cheese melts at around 40 °C. The latent heat of fusion of milk fat $\Delta H_{fus} = 84$ kJ/kg.

There are, in fact, two factors involved: the heat capacity (how much energy is stored in the materials and is available to be transferred to your cheek), and the thermal conductivity (how rapidly the heat energy can be transferred.) Since the heat energy is stored as random molecular kinetic motion, these are related. The key difference between crust, cheese, and sauce is the *water content*, which will generally tend to dominate the heat capacity. In other words, the higher the water content, the greater amount of heat that can be stored for the same rise in temperature. This is enhanced when a phase change (*i.e.* the cheese melting) occurs, since the energy added or lost during the phase change does *not* change the temperature!

Solution:

Calculate the heat energy available in each ingredient that can be transferred to your cheek. You can either (a) assume an initial starting temperature (either 21 or 25°C would be reasonable) to compare the total amount of energy stored by the time the pizza is a uniform 150 °C, or (b) simply look at how much heat energy can be transferred *i.e.* $\Delta T = (150 - 37) = 113$ °C.

Since the temperature of the pizza is above the cheese melting point and that of your cheek below, include the amount of energy required to melt the cheese, *i.e.*:

$$q_{cheese} = q_{heat} + q_{melt} = sm\Delta T + (84 \times 20)$$

Note that units of kJ/kg are the same as J/g, so the conversion factors cancel! Ultimately, the most liquid component will transfer its heat the fastest, as this is accomplished by molecular collisions (refer back to molecular kinetic theory), so it is in fact the sauce that burns first, even though there is less of it. Having said that, the cheese will likely cause greater burning if you keep it in contact with your lip, tongue or cheek too long. So always have pop with your pizza!

The Water and Wine Problem

You have a glass of water and a glass of wine. Assume that both are pure, homogeneous substances. (If it helps, consider the wine to be pure ethanol!)

1. Transfer exactly one teaspoon from the glass of water to the glass of wine and mix thoroughly.
2. Transfer exactly one teaspoon of this contaminated wine to the glass of water and mix thoroughly.

Consider the amount of water in the glass of wine, and the amount of wine in the glass of water: Which of the following statements is *true*?

- a) The amount of water in the wine is greater than the amount of wine in the water
- b) The amount of wine in the water is greater than the amount of water in the wine
- c) **The amount of water in the wine is equal to the amount of wine in the water**

Solution:

Most people – including university faculty members – get this wrong, and many still don't believe the answer after seeing the proof until they've worked it for themselves. So this really is a challenging problem, and I'd save it for your most advanced (or most self-assured!) students. I would also note that I found the proof, as originally presented, extremely hard to follow and therefore difficult to accept. Here goes...

- (i) Start with two containers, (1) and (2), containing pure A (water) and pure B (wine), so that:

$$V_1 = V_A^\circ \text{ and } V_2 = V_B^\circ$$

- (ii) Let the volume transferred from A to B in the first step be $V_{1,2}$. We now have:

$$V_1' = V_A^\circ - V_{1,2} \text{ and } V_2' = V_B^\circ + V_{1,2}$$

- (iii) The volume transferred from (2) back into (1) contains a small fraction of A (ΔV_A) from $V_{1,2}$.

Both glasses now have their original liquid volumes; neither contains pure A or pure B:

$$V_1'' = V_A^\circ - V_{1,2} + V_{2,1} \text{ and } V_2'' = V_B^\circ + V_{1,2} - V_{2,1}$$

- (iv) How much A is now in B in glass (2)? $V_{A,B} = V_{1,2} - \Delta V_A \therefore V_{1,2} = V_{A,B} + \Delta V_A$

How much B is now in A in glass (1)? $V_{B,A} = V_{2,1} - \Delta V_A \therefore V_{2,1} = V_{B,A} + \Delta V_A$

- (v) But $V_{1,2} = V_{2,1}$ therefore equating and rearranging from (iv) gives:

$$V_{A,B} + \Delta V_A = V_{B,A} + \Delta V_A \therefore V_{A,B} = V_{B,A} \text{ (Q.E.D)}$$

The Xenon Fluoride Problem

A sample of a compound comprising only xenon and fluorine was confined to a bulb with a pressure of 24 torr. Hydrogen was added to the bulb until the pressure was 96 torr. Passage of an electric spark through the mixture produced Xe and HF. After the HF was removed by reaction with solid KOH, the final pressure of xenon and unreacted hydrogen in the bulb was 48 torr. What is the empirical formula of the xenon fluoride in the original sample?

In the article giving this problem, it was noted that many chemistry professors other than those routinely teaching 1st-year would figure out the formula (XeF_4) and only then realise that it was “just an empirical formula question”! It is completely possible to solve this without needing the temperature, volume, gas constant, and pressure conversion factors, as long as it is assumed that all pressures were measured under the same conditions. Try it...

Sources:

1. Ruth Ben-Zvi, Bat-Sheva Eylon and Judith Silberstein, "Is an atom of copper malleable?" *J. Chem. Ed.*, 1986, 63(1), 64-66.
2. J. Dudley Herron, "Piaget for Chemists: Explaining what good students cannot understand." *J. Chem. Ed.*, 1975, 52(3), 146-150.
3. Douglas R. Mulford and William R. Robinson, "An inventory for alternate conceptions among first-semester general chemistry students." *J. Chem. Ed.*, 2002, 79(6), 739-744.
4. Lillian Bird, "Logical reasoning ability and student performance in general chemistry." *J. Chem. Ed.*, **2010**, 87(5), 541-546.
5. Mary B. Nakhleh, "Are our students conceptual thinkers or algorithmic problem solvers?" *J. Chem. Ed.*, 1993, 70(1), 52-55.
6. Jan Meyer and Ray Land, "Threshold concepts and troublesome knowledge: Linkages to ways of thinking and practising within the disciplines." *ETL Occasional Report Number 4*, **2003**. Available on-line at <http://www.tlrp.org/dspace/handle/123456789/177>
7. David Perkins, "The Many Faces of Constructivism." *Educational Leadership*, **1999**, 57(3), 6-11. Available on-line at <http://www.scribd.com/doc/32920521/Perkins-The-Many-Faces-of-Constructivism>
8. Vanessa Kind, *Beyond Appearances: Students' misconceptions about basic chemical ideas*, 2nd ed., Royal Society of Chemistry, 2004. (Note: original edition published under author's prior name Vanessa Barker no longer seems to be available) On-line from http://www.rsc.org/images/Misconceptions_update_tcm18-188603.pdf
9. Jan H. F. Meyer and Ray Land (eds.), *Overcoming Barriers to Student Understanding: Threshold concepts and troublesome knowledge*, Routledge/Taylor & Francis Group, 2006. This book includes the previously cited articles by Meyer & Land and Perkins in expanded form as separate chapters.
10. George M. Bodner, "Problem solving: The difference between what we do and what we tell students to do." *U. Chem. Ed.*, 2003, 7, 37-45. Available on-line from <http://www.rsc.org/Publishing/Journals/RP/issues/index.asp>
11. George M. Bodner, "Strengthening conceptual connections in introductory chemistry courses." *Chem. Ed. Res. Practice*, 2007, 8(1), 93-100. Available on-line from <http://www.rsc.org/Publishing/Journals/RP/issues/index.asp>
12. *Chemical Education: Towards Research-based Practice*, John K. Gilbert, Onno De Jong, Rosário Justi, David F. Reagust and Jan H. van Driel (eds), Kluwer Academic Publishers, Dordrecht/Boston/London, 2002.
13. Vanessa Kind, *Beyond Appearances: Students' misconceptions about basic chemical ideas*, 2nd ed., Royal Society of Chemistry, 2004. (Note: original edition published under author's prior name Vanessa Barker no longer seems to be available) http://www.rsc.org/images/Misconceptions_update_tcm18-188603.pdf
14. Vincente Talanquer, "Commonsense chemistry: A model for understanding students' alternative conceptions." *J. Chem. Ed.*, **2006**, 83(5), 811-816.

15. Keith S. Taber and Alejandra Garcia Franco, "Intuitive thinking and learning chemistry." *Education in Chemistry*, 2009, 46(2), March. Available on-line at <http://www.rsc.org/Education/EiC/issues/2009March/Index.asp>
16. Keith E. Stanovich, "Rational and Irrational Thought: The Thinking that IQ Tests Miss", *Scientific American Mind*, 2009, Nov/Dec. issue, 34-39. <http://www.scientificamerican.com/sciammind/?contents=2009-11>
17. Students' and Teachers' Conceptions and Science Education, final version compiled by Reinders Duit, University of Kiel Institute for Science Education, 2009. (Includes all branches of secondary school science) Available for download from <http://www.ipn.uni-kiel.de/aktuell/stcse/>